arXiv:1502.06466v1 [math.GM] 26 Mar 2014

On vector field generated by the Hopf map S^3 on S^2

Valerii Dryuma*

Institute of Mathematics and Informatics, AS RM,

5 Academiei Street, 2028 Kishinev, Moldova,

e-mail: valdryum@gmail.com

Abstract

The examples of solutions of the system of differential equations generated by the Hopf map $S^3 \to S^2$ are constructed. Their properties are discussed.

1 Introduction

The subject of our consideration is the system of ODE's in E^3 -space [1]

$$\frac{d}{dt}x(t) = 8\,\frac{4\,zx - y\left(x^2+y^2+z^2\right) + 4\,y}{\left(x^2+y^2+z^2+4\right)^2},$$

$$\frac{d}{dt}y(t) = 8\,\frac{4\,zy + x\left(x^2+y^2+z^2\right) - 4\,x}{\left(x^2+y^2+z^2+4\right)^2},$$

$$\frac{d}{dt}z(t) = \frac{24\,x^2 + 24\,y^2 - 8\,z^2 - \left(x^2+y^2+z^2\right)^2 - 16}{\left(x^2+y^2+z^2+4\right)^2} \qquad (1)$$

associated with the Hopf map $S^3 \to S^2$ of three dimensional sphere with equation $z_1\bar{z}_1 + z_2\bar{z}_2 = 1$ on two -dimensional sphere S^2 considered as set of the points $[\frac{z_1}{z_2}, 1]$ of the complex space C^2.

As it was shown in [1] from the system (1) can be obtained the relation

$$\frac{d\arctan(y(t)/x(t))}{dt} = 8\frac{x^2+y^2+z^2-4}{(x^2+y^2+z^2+4)^2},$$

$$\frac{d(x(t)^2+y(t)^2)}{dt} = 64z\frac{x^2+y^2}{(x^2+y^2+z^2+4)^2} \qquad (2)$$

and with the help of which was made qualitative analysis of the behavior of integral curves of the system (1) .

Here we consider properties of solutions of the system [1] more details.

---

[*] Work supported in part by Grant 14-0100389 and DAAD, Yena, Germany (2013)

2 Transformation

Theorem 1 *In spherical system of coordinates* r, ϕ, ψ

$$x(t) = r(t)\cos(\phi(t))\sin(\psi(t)),\ \ y(t) = r(t)\sin(\phi(t))\sin(\psi(t)),\ \ z(t) = r(t)\cos(\psi(t))$$

the system of equations (1) takes the form

$$\frac{d}{dt}\phi(t) = -8\,\frac{(r(t))^2 - 4}{16 + (r(t))^4 + 8\,(r(t))^2},$$

$$\frac{d}{dt}r(t) = \frac{\left((r(t))^4 + 8\,(r(t))^2 - 64\,(r(t))^2(\sin(\psi(t)))^2 + 16\right)\cos(\psi(t))}{16 + (r(t))^4 + 8\,(r(t))^2},$$

$$\frac{d}{dt}\psi(t) = -\frac{\sin(\psi(t))\left((r(t))^4 + 40\,(r(t))^2 - 64\,(r(t))^2(\sin(\psi(t)))^2 + 16\right)}{r(t)\left(16 + (r(t))^4 + 8\,(r(t))^2\right)}. \tag{3}$$

Now from the system (3) we obtain the equation

$$\left(\frac{d}{dr}\psi(r)\right)r^5\cos(\psi(r)) - 56\left(\frac{d}{dr}\psi(r)\right)r^3\cos(\psi(r)) + 64\left(\frac{d}{dr}\psi(r)\right)r^3(\cos(\psi(r)))^3 +$$

$$+16\left(\frac{d}{dr}\psi(r)\right)r\cos(\psi(r)) + \sin(\psi(r))r^4 - 24\,\sin(\psi(r))r^2 +$$

$$+64\,\sin(\psi(r))r^2(\cos(\psi(r)))^2 + 16\,\sin(\psi(r)) = 0, \tag{4}$$

which after substitution

$$\psi(r) = \arcsin(\sqrt{H(r)}) \tag{5}$$

takes the form

$$\left(r^5 + 8\,r^3 - 64\,r^3H(r) + 16\,r\right)\frac{d}{dr}H(r) + 2\,H(r)r^4 + 80\,H(r)r^2 - 128\,(H(r))^2r^2 + 32\,H(r) = 0. \tag{6}$$

General solution of the equation (6) is determined from implicit relation

$$4\,C_1\,\mathit{BesselI}(0, 1/2\sqrt{H(r)}r) + C_1\,\mathit{BesselI}(0, 1/2\sqrt{H(r)}r)r^2 -$$

$$-8\,C_1\,\mathit{BesselI}(1, 1/2\sqrt{H(r)}r)\sqrt{H(r)}r + 4\,\mathit{BesselK}(0, -1/2\sqrt{H(r)}r) +$$

$$+\mathit{BesselK}(0, -1/2\sqrt{H(r)}r)r^2 - 8\,\mathit{BesselK}(1, -1/2\sqrt{H(r)}r)\sqrt{H(r)}r = 0, \tag{7}$$

where C_1 is arbitrary.

In result we obtain that the function $\psi(r)$ from (5) can be expressed through the function $H(r)$ and then with a help of the first equation of the system (3) can be studied properties of the function $\phi(r)$.

3 Associated p.d.e.

Let us consider partial differential equation of first order which determine first integral $\Phi(r,\phi,\psi) = C$ of the system (3)

$$\left(-\sin(\psi)r^4 + 24\,\sin(\psi)r^2 - 64\,\sin(\psi)r^2\left(\cos(\psi)\right)^2 - 16\,\sin(\psi)\right)\frac{\partial}{\partial\psi}\Phi(r,\phi,\psi)+$$

$$\left(-8\,r^3 + 32\,r\right)\frac{\partial}{\partial\phi}\Phi(r,\phi,\psi)+$$

$$+\left(\cos(\psi)r^5 - 56\,\cos(\psi)r^3 + 64\,\left(\cos(\psi)\right)^3 r^3 + 16\,\cos(\psi)r\right)\frac{\partial}{\partial r}\Phi(r,\phi,\psi) = 0. \tag{8}$$

After substitution of the form

$$\psi = \arcsin(\xi),\ \ \Phi(r,\phi,\psi) = E(r,\phi,\xi),$$

$$\frac{\partial}{\partial\psi}\Phi(r,\phi,\psi) = \left(\frac{\partial}{\partial\xi}E(r,\phi,\xi)\right)\cos(\psi)$$

and taking in consideration the condition

$$\cos(\psi) = \sqrt{1-\xi^2}$$

we obtain from (8) the equation

$$\left(\frac{\partial}{\partial r}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}r^5 + 8\,\left(\frac{\partial}{\partial r}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}r^3 - 64\,\left(\frac{\partial}{\partial r}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}r^3\xi^2+$$

$$+16\,\left(\frac{\partial}{\partial r}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}r - 8\,\left(\frac{\partial}{\partial\phi}E(r,\phi,\xi)\right)r^3 + 32\,\left(\frac{\partial}{\partial\phi}E(r,\phi,\xi)\right)r-$$

$$-\left(\frac{\partial}{\partial\xi}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}\xi\,r^4 - 40\,\left(\frac{\partial}{\partial\xi}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}\xi\,r^2+$$

$$+64\,\left(\frac{\partial}{\partial\xi}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}\xi^3r^2 - 16\,\left(\frac{\partial}{\partial\xi}E(r,\phi,\xi)\right)\sqrt{1-\xi^2}\xi = 0. \tag{9}$$

It has solution of the form $\Phi(r,\phi,\psi) = F_2(\phi) + H(r,\psi)$, where

$$\frac{d}{d\phi}F_2(\phi) = c_2$$

and the function $H(r,\psi)$ satisfies to the equation

$$\left(\frac{\partial}{\partial r}H(r,\psi)\right)\cos(\psi)r^5 - 8\,\left(\frac{\partial}{\partial r}H(r,\psi)\right)\cos(\psi)r^3 + 16\,\left(\frac{\partial}{\partial r}H(r,\psi)\right)r^3\cos(3\,\psi)+$$

$$+16\,\left(\frac{\partial}{\partial r}H(r,\psi)\right)\cos(\psi)r - 8\,c_2r^3 + 32\,c_2r - \left(\frac{\partial}{\partial\psi}H(r,\psi)\right)\sin(\psi)r^4 + 8\,\left(\frac{\partial}{\partial\psi}H(r,\psi)\right)\sin(\psi)r^2-$$

$$-16\,\left(\frac{\partial}{\partial\psi}H(r,\psi)\right)r^2\sin(3\,\psi) - 16\,\left(\frac{\partial}{\partial\psi}H(r,\psi)\right)\sin(\psi) = 0 \tag{10}$$

where c_2 is parameter.

3.1 The (u, v)-method of integration

To derive particular solutions of the partial nonlinear differential equation

$$F(x, y, f_x, f_y, f_{xx}, f_{xy}, f_{yy}, f_{xxx}, f_{xyy}, f_{xxy}, ..) = 0 \tag{11}$$

we use parametric presentation of the functions and variables

$$f(x, y) \to u(x, t), \quad y \to v(x, t), \quad f_x \to u_x - \frac{u_t}{v_t} v_x, \quad f_y \to \frac{u_t}{v_t},$$

$$f_{yy} \to \frac{(\frac{u_t}{v_t})_t}{v_t}, \; f_{xy} \to \frac{(u_x - \frac{u_t}{v_t} v_x)_t}{v_t}, \; f_{xx} \to (f_x)_x - \frac{(f_x)_t}{v_t} v_x \tag{12}$$

where variable t is considered as parameter [2].

Remark that conditions of compatibility $f_{xy} = f_{yx}$ are fulfilled at the such type of presentation.

In result instead of equation (11) one gets the relation between the new variables $u(x, t)$, $v(x, t)$ and their partial derivatives

$$\Psi(u, v, u_x, u_t, v_x, v_t...) = 0. \tag{13}$$

This value coincides with the initial equation under the condition that $v(x, t) = t$ and takes a more general view in the choice of $u(x, t) = F(\omega(x, t), \omega_t...)$ and $v(x, t) = \Phi(\omega(x, t), \omega_t...)$.

In some cases the integration of equation (13) is a more simple problem than the equation (11).

With help of solutions of equation (13) $u(x, t)$, $v(x, t)$ the solutions of initial equation (11) $f(x, y)$ are obtained by exclusion of the parameter t from the conditions

$$f(x, y) = u(x, t), \; y = v(x, t). \tag{14}$$

Next, we apply this method for integrating equations (10).

In result of transformation of the function $H(r, \psi)$ and its derivatives with accordance of the rules (12)

$$H(\xi, \psi) = u\left(\xi, \eta\right), \; r = v(\xi, \psi)$$

$$\frac{\partial}{\partial r} H(r, \psi) = \frac{\frac{\partial}{\partial \xi} u(\xi, \psi)}{\frac{\partial}{\partial \xi} v(\xi, \psi)}, \quad \frac{\partial}{\partial \psi} H(r, \psi) = \frac{\partial}{\partial \psi} u(\xi, \psi) - \frac{\left(\frac{\partial}{\partial \psi} v(\xi, \psi)\right) \frac{\partial}{\partial \xi} u(\xi, \psi)}{\frac{\partial}{\partial \xi} v(\xi, \psi)} \tag{15}$$

the equation (10) takes form of the relation between the functions $u(\xi, \psi)$ and $v(xi, \psi)$

$$\Big(8 \sin(\psi)\xi^2 - \sin(\psi)\xi^4 - 16\,\xi^2 \sin(3\,\psi) - 16 \sin(\psi)\Big) \frac{\partial}{\partial \psi} u(\xi, \psi) \frac{\partial}{\partial \xi} v(\xi, \psi) +$$

$$+ \Big(\sin(\psi)\xi^4 - 8 \sin(\psi)\xi^2 + 16\,\xi^2 \sin(3\,\psi) + 16 \sin(\psi)\Big) \left(\frac{\partial}{\partial \xi} u(\xi, \psi)\right) \frac{\partial}{\partial \psi} v(\xi, \psi) +$$

$$+ \Big(16 \cos(\psi)\xi - 8 \cos(\psi)\xi^3 + \cos(\psi)\xi^5 + 16\,\xi^3 \cos(3\,\psi)\Big) \frac{\partial}{\partial \xi} u(\xi, \psi) -$$

$$-8\,c_2\xi^3 \frac{\partial}{\partial \xi} v(\xi, \psi) + 32\,c_2\xi\,\frac{\partial}{\partial \xi} v(\xi, \psi) = 0, \tag{16}$$

where the variable ξ is considered as parameter.

From here we can obtain the p.d.e by various ways. One of them can be obtained after the substitution of the form

$$v(\xi,\psi)=\xi\,\frac{\partial}{\partial\xi}\rho(\xi,\psi)-\rho(\xi,\psi),\quad u(\xi,\psi)=\frac{\partial}{\partial\xi}\rho(\xi,\psi) \tag{17}$$

that leads to the equation

$$\begin{gathered}\left(\sin(\psi)\xi^4-8\,\sin(\psi)\xi^2+16\,\xi^2\sin(3\,\psi)+16\,\sin(\psi)\right)\frac{\partial}{\partial\psi}\rho(\xi,\psi)+\\ +\xi^6\cos(\psi)-8\,\xi^4\cos(\psi)+16\,\xi^4\cos(3\,\psi)+16\,\xi^2\cos(\psi)-8\,c_2\xi^3+32\,c_2\xi=0\end{gathered} \tag{18}$$

to the function $\rho(\xi,\psi)$.

Solution of this equation has the form

$$\begin{gathered}\rho(\xi,\psi)=\frac{\xi^5\ln(\chi)}{16+40\,\xi^2+\xi^4}+\\ +16\,\frac{\xi^3\ln(\xi^4\chi^4+2\,\xi^4\chi^2+\xi^4+40\,\xi^2\chi^4-176\,\xi^2\chi^2+40\,\xi^2+16\,\chi^4+32\,\chi^2+16)}{16+40\,\xi^2+\xi^4}+\\ +8\,\frac{\xi^3\ln(\chi)}{16+40\,\xi^2+\xi^4}-\xi\,\ln(\chi^2+1)+16\,\frac{\xi\,\ln(\chi)}{16+40\,\xi^2+\xi^4}-\\ -64\,\_c_2\xi^6\arctan(1/32\,\frac{2\,\chi^2\,(16+40\,\xi^2+\xi^4)+32-176\,\xi^2+2\,\xi^4}{\sqrt{16\,\xi^2-24\,\xi^4+\xi^6}})\times\\ \times\left(16+40\,\xi^2+\xi^4\right)^{-1}\frac{1}{\sqrt{16\,\xi^2-24\,\xi^4+\xi^6}}-\\ -8\,\frac{c_2\xi^4\ln(\chi)}{16+40\,\xi^2+\xi^4}+\\ +256\,c_2\xi^4\arctan(1/32\,\frac{2\,\chi^2\,(16+40\,\xi^2+\xi^4)+32-176\,\xi^2+2\,\xi^4}{\sqrt{16\,\xi^2-24\,\xi^4+\xi^6}})\times\\ \times\left(16+40\,\xi^2+\xi^4\right)^{-1}\frac{1}{\sqrt{16\,\xi^2-24\,\xi^4+\xi^6}}+\\ +32\,\frac{\_c_2\xi^2\ln(\chi)}{16+40\,\xi^2+\xi^4}+F_1(\xi),\end{gathered} \tag{19}$$

where $\chi=\tan(1/2\,\psi)$ and $F_1(\xi)$-is arbitrary.

On base of of solutions of the equation (19) can be find the functions $u(\xi,\psi)$ and $v(\xi,\psi)$ from (17) and then get solutions of the equation (10) in parametric form

$$H(r,\psi)=u(\xi,\psi),\quad r=v(\xi,\psi).$$

References:

1. Aminov Yu.A., *Geometriya vektornogo polya.* M.: Nauka. Gl.red.fiz.-mat.lit.,(1990), p.1–208.

2. Dryuma V.S., *The Riemann and Einsten-Weyl geometries in theory of differential equations, their applications and all that.* A.B.Shabat et all.(eds.), New Trends in Integrability and Partial Solvability, Kluwer Academic Publishers, Printed in the Netherlands , 2004, p.115–156.